

A new Lower Cretaceous bird from China and tooth reduction in early avian evolution

Zhonghe Zhou and Fucheng Zhang Zhiheng Li

Proc. R. Soc. B 2010 **277**, 219-227 first published online 8 July 2009
doi: 10.1098/rsjb.2009.0885

Supplementary data

["Data Supplement"](#)

<http://rsjb.royalsocietypublishing.org/content/suppl/2009/07/07/rsjb.2009.0885.DC1.html>

References

[This article cites 21 articles, 3 of which can be accessed free](#)

<http://rsjb.royalsocietypublishing.org/content/277/1679/219.full.html#ref-list-1>

[Article cited in:](#)

<http://rsjb.royalsocietypublishing.org/content/277/1679/219.full.html#related-urls>

Subject collections

Articles on similar topics can be found in the following collections

[taxonomy and systematics](#) (145 articles)

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

A new Lower Cretaceous bird from China and tooth reduction in early avian evolution

Zhonghe Zhou* and Fucheng Zhang Zhiheng Li

Key Laboratory of Evolutionary Systematics of Vertebrates, Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences, 142 Xi-Wai-Da-Jie, PO Box 643, Beijing 100044, People's Republic of China

A new avian genus and species, *Zhongjianornis yangi* gen. et sp. nov., is reported from the Lower Cretaceous lacustrine deposits of the Jiufotang Formation in Liaoning, northeast China. The new taxon is characterized by possessing the following combination of features: upper and lower jaws toothless, snout pointed, humerus with large and robust deltopectoral crest, second phalanx of the major manual digit longer than the first phalanx, unguals of the alular and major digits of similar length and significantly shorter than the corresponding penultimate phalanges, tibiotarsus slender and more than twice the length of the tarsometatarsus, and metatarsal IV longer than the other metatarsals. Phylogenetic analysis indicates that *Zhongjianornis* is phylogenetically basal to *Confuciusornis* and the dominant Mesozoic avian groups, Enantiornithes and Ornithurae, and therefore provides significant new information regarding the diversification of birds in the Early Cretaceous. It also represents the most basal bird that completely lacks teeth, suggesting that tooth loss was more common than expected in early avian evolution and that the avian beak appeared independently in several avian lineages, most probably as a response to selective pressure for weight reduction. Finally, the presence of a significantly enlarged humeral deltopectoral crest suggests that *Zhongjianornis* shares with other basal birds such as *Jeholornis*, *Sapeornis* and *Confuciusornis* a distinctive mode of adaptation for flight contrasting with that seen in more advanced birds, which instead possess an elongated sternum and a prominent keel.

Keywords: Lower Cretaceous; bird evolution; beak; China

1. INTRODUCTION

Over 30 genera of Early Cretaceous birds have been reported in the last two decades from the Lower Cretaceous of northeastern China (Chiappe 2007, Gao *et al.* 2008, Zhou *et al.* 2008, O'Connor *et al.* 2009), documenting a burst of avian diversification that followed the appearance of the earliest bird *Archaeopteryx*. These early birds occur in association with hundreds of exceptionally preserved feathered dinosaurs, early mammals, pterosaurs, amphibians, flowered plants and insects, which constitute the famous Jehol Biota (Benton *et al.* 2008; Zhang *et al.* 2008).

Recently, a relatively large bird was recovered from the Jiufotang Formation in Jianchang, western Liaoning. The new material is represented by a nearly complete and articulated skeleton. The new bird displays a combination of features that are unknown in any previously reported taxon; in particular, it represents the most basal avian that had completely lost teeth. The robust boomerang-shaped furcula and the large deltopectoral crest of the humerus are strongly reminiscent of *Confuciusornis*. The new discovery adds to our understanding of avian biodiversity in the Early Cretaceous.

2. SYSTEMATIC PALEONTOLOGY

Aves Linnaeus, 1758.

Zhongjianornis yangi gen. et sp. nov.
(See figures 1–5).

(a) Holotype: A nearly complete articulated skeleton with some poorly preserved feather imprints (Institute of Vertebrate Paleontology and Paleoanthropology collection, IVPP V15900).

(b) Etymology: The genus and species names both honour the late Professor Zhongjian Yang (Chung-Chien Young), father of Chinese vertebrate paleontology and founder of the Institute of Vertebrate Paleontology and Paleoanthropology.

(c) Locality and horizon: Jianchang, Liaoning, northeastern China; Early Cretaceous Jiufotang Formation (He *et al.* 2004).

(d) Diagnosis: A pigeon-sized bird (table 1), distinguished from other early birds by possessing the following combination of features: toothless, snout pointed, humerus with a large and robust deltopectoral crest that is as wide as the humeral shaft and more than one-third the total length of the humerus, wing unguals small and slightly curved, metatarsal IV longer than II and III.

3. DESCRIPTION

The holotype of *Zhongjianornis yangi* gen. et sp. nov. is a nearly complete articulated skeleton of an adult individual (figures 1 and 2). The skull is characterized by a pointed snout and toothless upper and lower jaws. The maxillary and nasal processes of the premaxilla form an

* Author for correspondence (zhonghe@yeah.net).

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rspb.2009.0885> or via <http://rspb.royalsocietypublishing.org>.

One contribution to a Special Issue 'Recent advances in Chinese palaeontology'.



Figure 1. Holotype of *Zhongjianornis yangi* gen. et sp. nov. (IVPP V15900). Scale bar, 5 cm.

angle of less than 30° . The two nasal bones are long and strap-like and seem to be slightly constricted in the middle part and to flare towards the posterior end. The frontal is large and domed. The posterior end of the maxillary process of the left premaxilla contacts the maxilla, which is massively built and shorter than the premaxilla. The maxilla articulates posteriorly with the long, rod-shaped jugal.

A single occipital condyle is evident. Two flanges extend laterally from the posterior end of the basiptyergoid process. A broad bone lying near the basiptyergoid process is tentatively identified as a pterygoid, and a pair of slender bones between the nasal and maxilla is tentatively identified as the palatines.

The right mandible is well exposed in medial view. The dentary is strongly forked, with a long, straight ventral process and a short dorsal process that is slightly curved and nearly half the total length of the mandible. The angular appears massive, while the surangular is more slender. A triangular bone that tapers both anteriorly and posteriorly is recognized as the splenial, and a pair of slender rod-like bones is identified as hyoid elements (figure 3).

The cervical vertebrae are ventrally exposed. The posterior cervicals seem to be more elongated than the anterior ones. Because of crushing, it is difficult to determine the type of central articulation. The thoracic vertebrae are incompletely preserved, and their number is difficult to determine. One of the thoracics, located near the synsacrum, can be seen to have an opisthocoealous centrum. The lateral surface of each thoracic centrum appears to be excavated.

The thoracic ribs appear to be slender. At least three of them display tightly associated uncinatous process, which are expanded proximally but taper distally (figure 4a). No gastralia have been recognized, but it is




Figure 2. Line drawing of the holotype of *Zhongjianornis yangi* gen. et sp. nov. (IVPP V15900). Abbreviations: ca, caudal vertebra; ce, cervical vertebra; co, coracoid; dI-1, first phalanx of alular digit; dII-1, first phalanx of major digit; dII-2, second phalanx of major digit; fe, femur; fi, fibula; fu, furcula; hu, humerus; il, ilium; is, ischium; mcI, alular metacarpal; mcII, major metacarpal; mcIII, minor metacarpal; mtI, metatarsal I; pd, pedal digits; pu, pubis; r, radius; sc, scapula; st, sternum; sy, synsacrum; ti, tibia; tmt, tarsometatarsus; tv, thoracic vertebra; u, ulna; up, uncinatous process of rib. Scale bar, 5 cm.

unclear whether they are genuinely absent or simply not preserved.

The synsacrum is completely fused and exposed in the ventral view (figure 5a). However, its posterior end is obscured by the pelvis, and the transverse process of the last sacral is partly exposed through the acetabular opening. The number of sacral vertebrae is estimated to be no greater than eight.

The caudal series is incompletely preserved, although five caudal vertebrae are preserved in articulation posterior to the pelvis. A short rod-like chevron is associated with the most cranial of these vertebrae. The caudals are short and small, with short, flat transverse processes and apparently procoelous intercentral articulations. Near the right femur are a few more isolated caudals. One of them appears to be larger than the others and might possibly represent the pygostyle. Based on the morphology of the preserved caudals, it is probable that *Zhongjianornis* has only a short caudal series, although the presence of a pygostyle is questionable (figure 5a).

The left scapula is well preserved and articulates with the coracoid at an acute angle, while the left scapula and coracoid are more or less covered by the furcula and right coracoid (figure 4b). The elongate scapula is nearly as long as the humerus, with a curved and distally tapering shaft. The coracoid is also elongated and is a




Figure 3. (a) Skull of the holotype of *Zhongjianornis yangi* gen. et sp. nov. (IVPP V15900). (b) Line drawing of the holotype skull of *Zhongjianornis yangi* gen. et sp. nov. (IVPP V15900). Abbreviations: an, angular; ar, articular; bs, basisphenoid; de, dentary; fr, frontal; ju, jugal; ma, maxilla; na, nasal; pa, parietal; ?pal, ?palatine; pm, premaxilla; ?pt, ?pterygoid; sa, surangular; sp, splenial bone. Scale bar, 1 cm.

strut-like bone bearing a blunt acromion process. In contrast to ornithurine birds and at least one enantiornithine, *Protopteryx* (Zhang & Zhou 2000), the coracoid appears to lack a distinct procoracoid process. The sternal margin of the coracoid appears concave, with a maximum width less than half the length of the coracoid. The proximal part of the bone seems to bear a narrow supracoracoidal foramen.

The furcula is incompletely preserved and appears to lack a hypocleidum. Based on the available material, it can be reconstructed as having a robust boomerang-like shape, resembling the furculae of *Archaeopteryx* (Elzanowski 2002; Mayr *et al.* 2005), confuciusornithids (Hou *et al.* 1995; Chiappe *et al.* 1999) and other very basal birds. The furcula lacks grooves on its dorsal surface.

A thin bone between the right humerus and radius can be identified as an incompletely preserved sternum (one of the paired sternal plates), lying in proximity to the distal ends of some thoracic ribs. Although the outline of the whole sternum cannot be precisely reconstructed, this bone seems to be relatively short and more or less rounded, in contrast to elongated sterna of ornithurines. The sternum of *Zhongjianornis* also seems to lack a distinctly developed keel.

The forelimbs are exposed in dorsal view, and preserved in nearly complete articulation. The length ratio of the humerus + ulna + major metacarpal to the femur + tibiotarsus + metatarsal III is about 1.2, which is most similar to that of *Jeholornis* among known Mesozoic avians (Zhou & Zhang 2002). The humerus is robust, with a large deltopectoral crest that extends more than one-third the total length of the humerus. The deltopectoral crest is nearly as wide as the humeral shaft, with a rounded anterioproximal margin and a straight distal margin. The deltopectoral crest lacks a fenestra, in contrast to *Sapeornis* (Zhou & Zhang 2003) and confuciusornithids. The humeri of *Zhongjianornis* are straight and less twisted than in many more advanced birds. Proximally, the bicipital crest is prominent. The head of the humerus is not well exposed in either the right or the left forelimb, but seems not to be dorsally prominent. Distally, the humerus is slightly flared.

The ulna and radius are slightly longer than the humerus. The ulna is bow-shaped and is wider and more strongly curved than the radius. Distally the ulna seems to possess a semilunar external condyle. The right ulnare is preserved in association with the ulna and shows a distinctive metacarpal incision.




Figure 4. Close-up photos of *Zhongjianornis yangi* gen. et sp. nov. (IVPP V15900), showing (a) the thoracic ribs and uncinata processes, (b) the pectoral girdle, and (c) the hand. Abbreviations: dI-2, second phalanx of the alular digit; dII-3, third phalanx of the major digit; dIII-1, first phalanx of the minor digit; ul, ulnare; see figure 2 for other abbreviations. Scale bar, 1 cm.

The carpometacarpus is only fused at the proximal end, where a prominent carpal trochlea is present. The alular metacarpal is not fused with the major metacarpal and is a slender bone lacking an extensor process. The minor metacarpal is only about half as wide as the major metacarpal. It is only slightly curved near its distal end and does not fuse distally with the major metacarpal although the two bones are closely appressed. The major and minor metacarpals extend distally to about the same level, in contrast to enantiornithines, in which the minor metacarpal extends further distally.

The phalangeal formula of the manus is unclear. The alular digit is slender and does not extend to the distal end of the major metacarpal, and the first phalanx is much longer than the second (the unguis). The major digit comprises three phalanges. The first phalanx is the most robust of all the manual phalanges, whereas the second phalanx is shorter and more slender than the first phalanx and the third phalanx (the unguis) is shorter still. The minor digit is not preserved on the right wing, and only the first phalanx seems to be preserved on the

left. Thus, the phalangeal count of the minor digit is unclear. The unguis of the alular and major digits are only slightly curved and are about equal in length, both of them being significantly shorter than the non-ungual phalanges of the manus (figure 4c).

The right pelvis is exposed in the lateral view (figure 5a). Both ilia are preserved, with the right ilium overlapping much of the left. The ilium is mainly dorsoventrally extended, and its pre-acetabular portion is greater than its slightly tapering post-acetabular portion. The ischium is robust and curved posterodorsally, with a prominent ascending process that contacts the posterior end of the ilium. The posteriorly directed right pubis is preserved in articulation with the ilium. The rod-shaped shaft of the pubis is nearly straight along much of its proximal portion. The distal tip of the pubis is missing, but part of the pubic symphysis is preserved, although its length cannot be reconstructed. The proximal ischium and pubis are preserved in articulation with the ilium, forming a rounded acetabular foramen. No anti-trochanter is evident.




Figure 5. Close-up photos of *Zhongjianornis yangi* gen. et sp. nov. (IVPP V15900), showing (a) the pelvis and (b) the right foot. Abbreviations: mtII, metatarsal II; mtIII, metatarsal III; mtIV, metatarsal IV; pdI, pedal I; pdII, pedal II; pdIV, pedal IV; see figure 2 for other abbreviations. Scale bar, 1 cm.

The hind-limb elements are preserved in almost complete articulation. The femur is robust and nearly straight. It lacks a distinct neck but has a prominent ball-shaped head, which lacks a capital fossa. The femur is about two-thirds the length of the tibiotarsus and is much longer than the tarsometatarsus.

The tibiotarsus is long and slender, with the proximal and distal ends more or less crushed. Proximally, the tibiotarsus does not appear to form a cnemial crest. Distally, the tibiotarsus is well fused, but both the internal and external condyles are somewhat crushed and their shapes cannot be readily determined. The needle-shaped fibula is not fused with the tibiotarsus and extends along about three-quarters of the length of the latter bone. The right tibiotarsus appears shorter than the left, but we suspect that this is an artificial condition resulting from faulty reconstruction of the specimen by the local farmer before the specimen was obtained by us. Part of the middle segment was probably missing, as can be judged from a crack at the middle tibiotarsus.

The tarsometatarsus is relatively short, measuring only about 0.39 times the length of the tibiotarsus, which distinguishes it from any other known early birds (table 2). The right tarsometatarsus is completely preserved, but the left one is missing its distal end. The tarsometatarsus appears to be well fused at its proximal end. Based on the admittedly imperfect distal portion of the right tarsometatarsus, metatarsal IV appears to be longer than II and III (figure 5b), in contrast to most other early birds. However, we are less certain about the relative widths of the three major metatarsals. Metatarsal I is short, with an expanded distal end and a pointed proximal end. It appears to articulate with a facet on metatarsal II. Metatarsal V is absent. The pedal digits are nearly completely preserved, although they are greatly compressed. Digit I is moderately long, with the ungual exceeding the first phalanx in length. Digit II has a short first phalanx and a relatively long second phalanx, which, however, is shorter than the ungual. Digit III is slightly wider than the other digits. The non-ungual phalanges of digit IV are subequal in length and relatively short compared with those of the other digits. All of the pedal unguals are long and curved.

4. COMPARISON AND DISCUSSION

The new specimen represents a bird that is relatively large compared with other known birds from the Lower Cretaceous (table 1). It can be most readily distinguished from other taxa by the following features: skull with a pointed snout and toothless jaws, and robust humerus with a distinctively enlarged deltopectoral crest. Among Early Cretaceous birds of comparable or greater size, both *Jeholornis* and *Sapeornis* have a greater forelimb to hind-limb length ratio (Zhou & Zhang 2003). *Jeholornis* also has a unique skull adapted for seed-eating, with robust jaws bearing reduced teeth (Zhou & Zhang 2002). The skull of *Sapeornis* is adapted for herbivory, but still retains large teeth on the premaxilla, and the humerus of this taxon differs from that of *Zhongjianornis* in displaying a distinct fenestra on the deltopectoral crest. Furthermore, the femur and metatarsal III of *Zhongjianornis* are both significantly shorter in comparison to the tibiotarsus than those in *Sapeornis*.

The confuciusornithids are a group of birds from northeast China that are similar in size to *Zhongjianornis*, and they also resemble this taxon in having completely toothless jaws (Zhang *et al.* 2008). However, *Zhongjianornis* can be easily distinguished from confuciusornithids by its pointed snout and by the small size of the ungual of the alular manual digit (Hou *et al.* 1995; Chiappe *et al.* 1999). Furthermore, *Zhongjianornis* has a greater forelimb to hind-limb ratio than is seen in confuciusornithids (table 2).

Finally, the basal ornithurines *Hongshanornis* (Zhou & Zhang 2005) and *Archaeorhynchus* (Zhou & Zhang 2006) are two additional Chinese early birds that completely lack teeth. However, both are smaller than *Zhongjianornis*. *Hongshanornis* has far more slender limb bones, lacks an extremely large deltopectoral crest and displays a much smaller forelimb to hind-limb ratio. *Archaeorhynchus*, by contrast, is readily distinguishable from *Zhongjianornis* by its greater forelimb to hind-limb ratio. More advanced ornithurines such as *Yixianornis* and *Yanornis* (Zhou & Zhang 2001; Clarke *et al.* 2006) are toothed. All known enantiornithines from the Early Cretaceous of China have retained at least some teeth in the skull

Table 1. Measurements of *Zhongjianornis yangi* gen. et sp. nov. (IVPP V15900).

elements	measurements (mm)
skull length	61
synsacrum length	28 ^a
synsacrum, maximum width	8
scapula length (right)	59
coracoid length (right)	36
coracoid, distal maximum width (left)	16
humerus length (left)	71
humerus, midshaft width (left)	6
ulna length (right)	74
ulna, midshaft width (right)	5
radius length (right)	73 ^a
radius, midshaft width (right)	3.5
carpometacarpus length (right)	34
major metacarpal (right)	31
major metacarpal, midshaft width (right)	3.5
minor metacarpal, midshaft width (right)	2
alular metacarpal length (right)	8
major digit phalanx-1 length (left)	18
major digit phalanx-2 length (left)	19.5
major digit phalanx-3 length (ungual, right)	5 ^b
alular digit phalanx-1 length (right)	15
alular digit phalanx-2 length (ungual, right)	5.5 ^b
ilium length (left)	39
ischium length (right)	25 ^b
pubis length (right)	32 ^b
femur length (left)	46.5
femur length (right)	48
tibiotarsus length (left)	65 ^a
tibiotarsus length (right)	71
fibula length (right)	53
tarsometatarsus length (left)	29 ^a
tarsometatarsus length (right)	27 ^b
metatarsal I length (left)	6
metatarsal II length (left)	26
metatarsal III length (left)	28 ^a
metatarsal IV length (left)	29 ^a
pedal digit I-1 length (left)	5
pedal digit I-2 length (ungual, left)	8 ^b
pedal digit II-1 length (left)	6.5
pedal digit II-2 length (left)	8
pedal digit II-3 length (ungual, left)	11
pedal digit III-1 length (left)	6.5
pedal digit III-2 length (left)	7 ^a
pedal digit III-3 length (left)	7 ^a
pedal digit III-4 length (ungual, left)	12
pedal digit IV-1 length (right)	7
pedal digit IV-2 length (right)	5
pedal digit IV-3 length (right)	4 ^a
pedal digit IV-4 length (right)	6.5
pedal digit IV-5 length (ungual, right)	9.5 ^b

^aEstimated value.^bPreserved length.

(Zhang & Zhou 2000; Zhou *et al.* 2005) and are often much smaller than *Zhongjianornis*. This is true even of the largest enantiornithine, *Pengornis* (Zhou *et al.* 2008), which also has greater forelimb to hind-limb and femur to tibiotarsus ratios than *Zhongjianornis* (table 2).

Phylogenetic analysis indicates that *Zhongjianornis* represents one of the most basal known birds. This taxon is more derived than *Archaeopteryx*, *Sapeornis* and

Table 2. Measurements (mm) and proportions of selected limb elements of *Zhongjianornis yangi* (IVPP V15900) and some other early birds from China. l. and r. indicate left and right sides, respectively. Abbreviations are as in figures 2 and 5.

	<i>Zhongjianornis yangi</i> (IVPP V15900)	<i>Jeholornis prima</i> (IVPP V13274)	<i>Sapeornis chaoyangensis</i> (IVPP V12698)	<i>Confuciusornis sanctus</i> (IVPP V11619)	<i>Pengornis houi</i> (IVPP V15336)	<i>Longipteryx chaoyangensis</i> (IVPP V12325)	<i>Archaeorhynchus spathula</i> (IVPP V14287)	<i>Hongshanornis longicresta</i> (IVPP V14533)	<i>Yixianornis grabaui</i> (IVPP V12631)
humerus	71 (l.)	110 (r.)	127 (l.)	52 (r.)	64 (l.)	45 (r.)	53 (l.)	26 (l.)	49 (r.)
ulna	74 (r.)	109 (r.)	133 (l.)	47 (r.)	71 (l.)	47 (r.)	56 (l.)	24 (l.)	50 (r.)
major metacarpal	31 (r.)	47 (r.)	57 (l.)	27 (r.)	27 (r.)	19 (r.)	25 (l.)	13 (l.)	21 (r.)
femur	48 (r.)	75 (r.)	80 (l.)	47 (l.)	48 (r.)	31 (r.)	37 (l.)	22 (l.)	42 (r.)
tibiotarsus	71 (r.)	88 (r.)	84 (l.)	54 (l.)	50 (r.)	32 (r.)	42 (l.)	38 (l.)	54 (r.)
metatarsal III	28 (l.)	47 (r.)	44 (l.)	25 (l.)	26 (l.)	21 (r.)	20 (l.)	22 (l.)	26 (r.)
femur/tibiotarsus	0.68	0.85	0.95	0.87	0.96	0.97	0.88	0.58	0.78
mtIII/ti	0.39	0.53	0.52	0.46	0.52	0.66	0.48	0.58	0.48
hu + u + mcII/fe + ti + mtIII	1.20	1.26	1.52	1.0	1.35	1.32	1.35	0.79	0.98




Figure 6. Phylogeny of *Zhongjianornis yangi* gen. et sp. nov. Phylogenetic analysis was conducted using NONA 2.0 and WINCLADA. All characters were unordered. A total of 205 morphological characters were used, following Zhou *et al.* (2008). The data matrix from Zhou *et al.* (2008) was used in a revised form. Tree length = 487; CI = 0.55; RI = 0.77. The cladogram is based on the strict consensus of four most parsimonious trees.

Jeholornis, but is more basal than other known taxa including *Confuciusornis* and the two dominant Mesozoic avian groups, the enantiornithines and ornithurines (figure 6). The new discovery adds to the known diversity of Lower Cretaceous avian assemblages. Furthermore, it is notable that the most basal currently known avian taxa are all relatively large in comparison with the more derived enantiornithines, confirming that the evolution of improved powered flight was coupled with a size decrease on the lineage leading to enantiornithines. The new cladogram is generally consistent with a recent analysis by Zhou *et al.* (2008), except that *Sapeornis* occupies a more basal position than *Jeholornis* in the current analysis. This result is surprising, as it contrasts with all previous analyses. The new topology readily explains the presence of a primitive non-strut-like coracoid in *Sapeornis*, but would also suggest that the long skeletal tail in *Jeholornis* could have evolved secondarily.

Among previously reported beaked birds from the Mesozoic of China, confuciusornithids represent the most basal forms to display this advanced feature. *Jeholornis* and *Sapeornis* have greatly reduced their teeth, but in neither of these taxa are teeth completely lacking (Zhou *et al.* 2005; Zhou & Zhang 2007). All known Chinese enantiornithines are also toothed, although some of them, including *Longipteryx* (Zhang *et al.* 2001), *Longirostravis* (Hou *et al.* 2004) and *Shanweimiao* (O'Connor *et al.* 2009), have their teeth restricted to the premaxilla and anterior dentary. Among known basal ornithurines, both *Hongshanornis* and *Archaeorhynchus* are completely toothless. An analysis of 31 avian genera from the Lower Cretaceous of China shows that seven of them are edentulous (table 3), and most are either completely toothed (i.e. teeth present in premaxilla,

maxilla and dentary) or partially toothed (i.e. teeth absent on either one or two of the same three bones). The avian assemblage from the Lower Cretaceous of China documents a transition from completely toothed birds in the Late Jurassic to completely toothless ones in the Cenozoic.

The new bird also represents the most basal form to completely lose its teeth, although tooth loss undoubtedly occurred independently in several early avian lineages. Among known toothless birds, *Confuciusornis* is the next most advanced. We assume that tooth loss in both cases was probably owing to selective pressure for a reduction in body weight, and it would be especially advantageous to reduce the weight of the head rather than, for instance, the trunk, because the former is further from the centre of gravity. Many ornithurines that retained teeth, such as *Yanornis* and *Yixianornis*, also appear to have been strong flyers, and the selection pressure for weight reduction was probably less severe in these taxa.

The great variations in jaw and rostral morphology that existed among early birds may also suggest that significant trophic differentiation played a key role in the explosive radiation of birds approximately 20 million years after *Archaeopteryx* (Chiappe & Walker 2002; Chiappe 2007; Zhou & Zhang 2007).

Zhongjianornis appears to retain a boomerang-shaped furcula like those of more basal avians and non-avian theropods. Unlike *Archaeopteryx*, *Jeholornis* and *Sapeornis*, it possesses well-developed uncinata processes, as in *Confuciusornis* and more derived taxa.

The humerus has a robust deltopectoral crest, much as in confuciusornithids, but lacks a fenestra of the kind seen in *Sapeornis* and most confuciusornithids. Zhang *et al.* (2008) showed that a remarkably enlarged humeral

Table 3. List of 31 known avian genera from the Lower Cretaceous of China.

	genera
completely toothed (11)	<i>Protopteryx</i> , <i>Eoenantiornis</i> , <i>Cathayornis</i> , <i>Sinornis</i> , <i>Liaoxiornis</i> , <i>Fibeimia</i> , <i>Vescornis</i> , <i>Songlingornis</i> , <i>Yanornis</i> , <i>Yixianornis</i> , <i>Pengornis</i>
partially toothed (6)	<i>Jeholornis</i> , <i>Sapeornis</i> , <i>Boluochia</i> , <i>Longirostravis</i> , <i>Longipteryx</i> , <i>Shanweimiao</i>
edentulous (7)	<i>Confuciusornis</i> , <i>Changchengornis</i> , <i>Jinzhourornis</i> , <i>Eoconfuciusornis</i> , <i>Hongshanornis</i> , <i>Archaeorhynchus</i> , <i>Zhongjianornis</i>
unknown (7)	<i>Gansus</i> , <i>Chaoyangia</i> , <i>Eocathayornis</i> , <i>Otogornis</i> , <i>Liaoningornis</i> , <i>Paraprotopteryx</i> , <i>Zhongornis</i>

deltpectoral crest represents an adaptation for increasing flight capability between 131 and 120 Ma for confuciusornithids. It is notable that both *Jeholornis* and *Sapeornis* have a large deltopectoral crest, a condition unknown in enantiornithine and ornithurine birds (Zhou & Zhang 2001; Clarke & Norell 2002; Clarke *et al.* 2006). Besides, none of these most basal birds possesses an elongated sternum or a prominent sternal keel. The presence of a large deltopectoral crest in these basal birds probably compensated to some extent for the lack of an elongate, keeled sternum.

The proportions of the hind-limb and pes, as well as the shapes of the pedal unguals, seem to indicate an arboreal habit for *Zhongjianornis*. For instance, metatarsal III is only about 0.39 times as long as the tibiotarsus, a smaller ratio than in other known Early Cretaceous birds. The ratio of the length of the femur to that of the tibiotarsus is 0.68, also a smaller value than is typical of Early Cretaceous birds. The pedal unguals are all long and curved, and significantly longer than other phalanges. The hand of *Zhongjianornis* still retains some primitive features, such as the presence of a long major digit in which the second phalanx is longer than the first. However, it is also notable that the wing unguals seem to be relatively reduced and less curved than in more basal birds such as *Jeholornis*, *Sapeornis* and *Confuciusornis*, possibly indicating that they have lost their former presumed role in climbing.

We are grateful to Corwin Sullivan for reading the manuscript and providing many suggestions. We also thank Xiaolin Wang for help in the field, Yan Li for fossil preparation and Wei Gao for photos. This research was funded by the Major Basic Research Projects (2006CB806400) of MST of China and the National Natural Science Foundation of China (40121202).

REFERENCES

Benton, M. J., Zhou, Z., Orr, P. J., Zhang, F. & Kearns, S. L. 2008 The remarkable fossils from the Early Cretaceous Jehol Biota of China and how they have changed our knowledge. *Proc. Geol. Assoc.* **119**, 209–228. (doi:10.1016/S0016-7878(08)80302-1)

Chiappe, L. M. 2007 *Glorified dinosaurs: the origin and early evolution of birds*. Hoboken, NJ: John Wiley & Sons, Inc.

Chiappe, L. M. & Walker, C. A. 2002 Skeletal morphology and systematics of the Cretaceous Euenantiornithes (Ornithothoraces: Enantiornithes). In *Mesozoic birds: above the heads of dinosaurs* (eds L. M. Chiappe & L. M. Witmer), pp. 240–267. Berkeley, CA: University of California Press.

Chiappe, L. M., Ji, S., Ji, Q. & Norell, M. A. 1999 Anatomy and systematics of the Confuciusornithidae (Theropoda: Aves) from the Late Mesozoic of Northeastern China. *Bull. Am. Mus. Natl Hist.* **242**, 1–89.

Clarke, J. A. & Norell, M. A. 2002 The morphology and phylogenetic position of *Apsaravis ukhaana* from the Late Cretaceous of Mongolia. *Am. Mus. Novit.* **3387**, 1–46. (doi:10.1206/0003-0082(2002)387<0001:TMAPPO>2.0.CO;2)

Clarke, J., Zhou, Z. & Zhang, F. 2006 Insight into the evolution of avian flight from a new clade of Early Cretaceous ornithurines from China and the morphology of *Yixianornis grabaui*. *J. Anat.* **208**, 287–308. (doi:10.1111/j.1469-7580.2006.00534.x)

Elzanowski, A. 2002 Archaeopterygidae (Upper Jurassic of Germany). In *Mesozoic birds: above the heads of dinosaurs* (eds L. M. Chiappe & L. M. Witmer), pp. 129–159. Berkeley, CA: University of California Press.

Gao, C., Chiappe, L. M., Meng, Q., O'Connor, K., Wang, X., Cheng, C. & Liu, J. 2008 A new basal lineage of Early Cretaceous birds from China and its implications on the evolution of the avian tail. *Palaeontology* **51**, 775–791. (doi:10.1111/j.1475-4983.2008.00793.x)

He, H., Wang, X., Zhou, Z., Wang, F., Boven, A., Shi, G. & Zhu, R. 2004 Timing of the Jiufotang Formation (Jehol Group) in Liaoning, northeastern China and its implications. *Geophys. Res. Lett.* **31**, L12605. (doi:10.1029/2004GL019790)

Hou, L., Zhou, Z., Martin, L. D. & Feduccia, A. 1995 A beaked bird from the Jurassic of China. *Nature* **377**, 616–618. (doi:10.1038/377616a0)

Hou, L., Chiappe, L. M., Zhang, F. & Chuong, C. 2004 New Early Cretaceous fossil from China documents a novel trophic specialization for Mesozoic birds. *Naturwissenschaften* **91**, 22–25. (doi:10.1007/s00114-003-0489-1)

Mayr, G., Pohl, B. & Peters, D. S. 2005 A well-preserved *Archaeopteryx* specimen with theropod features. *Science* **310**, 1483–1486. (doi:10.1126/science.1120331)

O'Connor, J. K., Wang, X., Chiappe, L. M., Gao, C., Meng, Q., Cheng, X. & Liu, J. 2009 Phylogenetic support for a specialized clade of Cretaceous enantiornithine birds with information from a new species. *J. Vert. Paleontol.* **29**, 188–204.

Zhang, F. & Zhou, Z. 2000 A primitive enantiornithine bird and the origin of feathers. *Science* **290**, 1955–1959. (doi:10.1126/science.290.5498.1955)

Zhang, F., Zhou, Z., Hou, L. & Gu, G. 2001 Early diversification of birds: evidence from a new opposite bird. *Chin. Sci. Bull.* **46**, 945–950. (doi:10.1007/BF02900473)

Zhang, F., Zhou, Z. & Benton, M. J. 2008 A primitive confuciusornithid bird from China and its implications for early avian flight. *Sci. China (D)* **51**, 625–639.

Zhou, Z. & Zhang, F. 2001 Two new ornithurine birds from the Early Cretaceous of western Liaoning, China. *Chin. Sci. Bull.* **46**, 1258–1264.

Zhou, Z. & Zhang, F. 2002 A long-tailed, seed-eating bird from the Early Cretaceous of China. *Nature* **418**, 405–409. (doi:10.1038/nature00930)

- Zhou, Z. & Zhang, F. 2003 Anatomy of the primitive bird *Sapeornis chaoyangensis* from the Early Cretaceous of Liaoning, China. *Can. J. Earth Sci.* **40**, 731–747. (doi:10.1139/e03-011)
- Zhou, Z. & Zhang, F. 2005 Discovery of a new ornithurine bird and its implication for Early Cretaceous avian radiation. *Proc. Natl Acad. Sci. USA* **102**, 18 998–19 002. (doi:10.1073/pnas.0507106102)
- Zhou, Z. & Zhang, F. 2006 A beaked basal ornithurine bird (Aves, Ornithurae) from the Lower Cretaceous of China. *Zool. Script.* **35**, 363–373. (doi:10.1111/j.1463-6409.2006.00234.x)
- Zhou, Z. & Zhang, F. 2007 Mesozoic birds of China—a synoptic review. *Front. Biol. China* **2**, 1–14. (doi:10.1007/s11515-007-0001-y)
- Zhou, Z., Chiappe, L. M. & Zhang, F. 2005 Anatomy of the Early Cretaceous bird *Eoenantiornis buhleri* (Aves: Enantiornithes) from China. *Can. J. Earth Sci.* **42**, 1331–1338. (doi:10.1139/e05-038)
- Zhou, Z., Clarke, J. & Zhang, F. 2008 Insight into diversity, body size, and morphological evolution from the largest Early Cretaceous enantiornithine bird. *J. Anat.* **212**, 565–577. (doi:10.1111/j.1469-7580.2008.00880.x)