

Asian Paleocene charophyte records demonstrate Eocene dispersals from Asia to Europe

Wenxin Cao,^{1,2,3,4,‡} Sha Li,^{3,*} Qiang Li,^{1,2,5} Thomas A. Stidham,^{1,2,5} Xiaoqiao Wan,⁶ and Xijun Ni^{1,2,5,7*}

¹Key Laboratory of Vertebrate Evolution and Human Origins, Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences, Beijing, 100044, China <caowenxin@ivpp.ac.cn>, <liqiang@ivpp.ac.cn>, <presbyornis@gmail.com>, <nixijun@ivpp.ac.cn>

²CAS Center for Excellence in Life and Paleoenvironment, Beijing 100044, China

³State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology and Palaeontology, Centre for Excellence in Life and Paleoenvironment, Chinese Academy of Sciences, 39 East Beijing Road, Nanjing 210008, China <shali@nigpas.ac.cn>

⁴General Prospecting Institute of China National Administration of Coal Geology, Beijing 100039, China

⁵University of the Chinese Academy of Sciences, Beijing 100049, China

⁶State Key Laboratory of Biogeology and Environmental Geology, School of Earth Sciences and Resources, China University of Geosciences, Beijing 100083, China <wanxq@cugb.edu.cn>

⁷CAS Center for Excellence in Tibetan Plateau Earth Sciences, Beijing 100104, China

Abstract.—A latest Paleocene charophyte flora collected from the South Gobi area in the Junggar Basin, western China, includes the geographically widespread taxa *Peckichara torulosa* var. *varians* (Dollfus and Fritel, 1919) Sanjuan, Vicente, and Eaton, 2020, *Lychnothmanus vectensis* (Groves, 1926) Soulié-Märsche, 1989, and *Gyrogona lemani capitata* Grambast and Grambast-Fessard, 1981. *Lychnothmanus vectensis* (as *Lychnothmanus* aff. *L. vectensis*) is known from the Cretaceous–Paleocene transition in eastern China and the latest Paleocene in western China, with likely additional records from the United States (Utah). The earliest European records of *L. vectensis* are from the late Eocene to early Oligocene in Spain, France, and England. Similarly, the oldest record of *G. lemani capitata* is from the latest Paleocene in the South Gobi area, with younger records from the middle Eocene of France. These latest Paleocene gyrogonite assemblages demonstrate the origin of these charophyte lineages in Asia. The dispersal of these charophytes from Asia to Europe in the middle to late Eocene appears to have occurred before the retreat of the Turgai Strait in both the Tarim area and the Siberian Basin by the end of the late Eocene and before the “Grande Coupure” in Europe and the Mongolian Remodelling in Asia during the Eocene–Oligocene transition. We hypothesize that waterbirds may have facilitated this intercontinental dispersal, and that idea is supported by the shared occurrence of avian groups in Central Asia and Europe in the middle and late Eocene.

Introduction

Charophyte paleobiogeography has been studied widely in the past two decades. Bhatia (2006) suggested that the living Eurasian species *Lychnothamnus barbatus* (Meyen, 1827) von Leonhardi, 1863 dispersed from Western Europe (late Miocene) eastward to other parts of Europe and Central Asia in the Pliocene, with hypothesized dispersals to the Himalayas in the late Pliocene and to Australia in the Pleistocene. Martín-Closas and Wang (2008) studied the paleobiogeography of the well-documented, long-lasting fossil record of Cretaceous charophyte *Atopochara trivolvus* Peck, 1941 and concluded that the species probably originated in Eastern Europe, later achieving a global distribution. Some charophyte species have dispersed much faster than others. For example, Sanjuan and Martín-Closas (2015a) studied the Eocene–Oligocene *Nitellopsis* (*Tectochara*) *merianii* Braun and Unger in Unger, 1852

comb. Grambast and Soulié-Märsche, 1972–*Nitellopsis obtusa* (Desvaux in Loiseleur, 1810) Groves, 1919 and *Lychnothamnus stockmansii* (Grambast, 1957) Soulié-Märsche, 1989–*Lychnothamnus major* (Grambast and Paul, 1965) Soulié-Märsche, 1989 lineages, and they surmised that the *L. stockmansii*–*L. major* lineage achieved a Eurasian distribution soon after its first appearance in the late Eocene, and the *N. (T.) merianii*–*N. obtusa* lineage expanded to Eastern Europe from Spain about 10 million years after its first appearance.

From a paleogeographical perspective, Europe and Western Asia were separated from Northeastern Asia by the Turgai Straits from the Middle Jurassic (Martin and Averianov, 2004; Joyce and Rabi, 2015) to the end of the late Eocene (McKenna, 1975; Martin and Averianov, 2004; Bosboom et al., 2017). The Turgai Strait, also known as the West Siberian Sea, was a large shallow saline water body that connected the Arctic Ocean and Tethys Sea and separated the Asian and European continents. The dispersal of charophytes from Asia to Europe in the middle to late Eocene appears to have occurred before the retreat of the Turgai Strait in both the Tarim area and the Siberian Basin by the end of the late Eocene. Our recovery of late Paleocene

[‡]Wenxin Cao and Sha Li contributed equally to this work.

*Corresponding author.




Figure 1. Geographic location. Maps showing location of the South Gobi area in the Northeastern Junggar Basin in Xinjiang, China.

charophyte fossils in Central Asia (western China) adds greatly to the paleobiogeographic story of intercontinental dispersals and distributions of charophyte taxa in the Paleogene. We use charophyte data to discuss potential agents of dispersal and paleoenvironments and to form the basis to explain the expansion and colonization of various areas by charophytes.

Geological background

The Junggar Basin is one of the largest sedimentary basins in Xinjiang, northwestern China. It is surrounded by the Zaier and Hala' alate Mountains to the northwest, the Aletay and Kelamaili Mountains to the northeast, and the Tian Shan and Bogda mountains to the south (Tang et al., 2020). The basin includes many fossiliferous horizons from the Cretaceous and Paleogene. The latest Paleocene South Gobi fossil area (46°5'19"N, 88°11'9"E) is situated in an unnamed early Cenozoic lithological unit located in the northeastern part of the Junggar Basin in the Gurbantünggüt Desert (Fig. 1). This horizon is known primarily for its fossil vertebrate assemblage (Ni et al., 2016), but fossils of charophytes also have been discovered in this area. This fossiliferous unnamed geological unit comprises a set of sediments consisting of reddish-brown mudstones interbedded with a few grayish-green or yellowish-gray fluvial sandstone beds or lenses (Ni et al., 2016). The fossils of charophytes were recovered from the sandy mudstone unit (at the height of ~7.5 m in Fig. 2).

The geological age of the deposit was determined through mammalian biostratigraphy (Ni et al., 2016). Mammalian fossils in the unit include species of insectivorans (*Bumbanius ningi* Missiaen and Smith, 2008 and *Asionyctia guoi* Missiaen and Smith, 2005), Glires (*Tribosphenomys minutus* Meng and Wyss, 2001 and *Neimengomys qii* Meng et al., 2007), plesiadapiforms (*Subengius mengi* Smith, Van Itterbeeck, and Missiaen, 2004), arctostyloids (*Anatolestylops* sp. aff. *Anatolestylops*

zhaii Wang et al., 2008), deltatheroidan marsupials (*Gurbantodonta kara* Ni et al., 2016), and multituberculates (*Mesodmops* cf. *M. tenuis* Missiaen and Smith, 2008). Among them, seven species (all except *Anatolestylops* sp. aff. *A. zhaii*) were reported as also occurring in the well-constrained late Paleocene Subeng Fauna from the Erlian Basin of Inner Mongolia, China (Ni et al., 2016). In addition, the specimens of *Anatolestylops* sp. aff. *A. zhaii* (an arctostyloid) from the South Gobi Fauna are much larger than those of *Palaeostylops iturus* Matthew and Granger, 1925 from the late Paleocene Subeng Fauna, and they approach the size of *Anatolestylops zhaii* from the early Eocene Nomogen III Fauna of Inner Mongolia. Therefore, the mammalian fossils and the South Gobi deposit are thought to be from the latter part of the Paleocene (lacking any Eocene biostratigraphic indicators). The Paleocene–Eocene Boundary in the Erlian Basin lies within or just below the “*Gomphos*” bed, and the absence of the common *Gomphos* (Glires, Mammalia) in the South Gobi area also points to an age older than the Eocene (Ni et al., 2016 and references therein) (Fig. 2). Therefore, this latest Paleocene South Gobi Fauna (and charophyte flora) can be correlated to the late Gashatan Asian Land Mammal Age, which is roughly equivalent to the Clarkforkian North American Land Mammal Age, with the age estimate of ~56–57 Ma (Fig. 2).

Materials and methods

Approximately 20 tons of sedimentary matrix (composing one massive sample) was collected from the latest Paleocene fossiliferous layer in Section A (Section, Fig. 1). The sample was screen washed with a minimum mesh size of 0.2 mm to maximize fossil recovery. For the semi-quantitative analyses, ~100 gyronite specimens were measured for each species (e.g., *Gyrogona lemani capitata* Grambast and Grambast-Fessard, 1981). When there were fewer than 100 specimens available for a species (e.g., *Lychnothmanus vectensis* (Groves, 1926) Soulié-Marsche, 1989 and *Peckichara torulosa* var. *varians* (Dollfus and Fritel, 1919) Sanjuan, Vicente, and Eaton, 2020), all the specimens were measured. Morphometric data from specimens of *G. lemani capitata* and *L. vectensis* were compiled into statistical charts, but charts were not made for *P. torulosa* var. *varians* because there are too few specimens. The specimens were measured using the ZEN 2 system stored with a Zeiss stereomicroscope (Stemim 508) at the Nanjing Institute of Geology and Paleontology, Chinese Academy of Sciences. Specimens were imaged using a Zeiss Evo MA25 scanning electron microscope at the Key Laboratory of Vertebrate Evolution and Human Origins of the Chinese Academy of Sciences, Beijing.

Repository and institutional abbreviation.—All specimens in this study are deposited in the Institute of Vertebrate Paleontology and Paleoanthropology (IVPP), Chinese Academy of Sciences, Beijing, China.

Systematic paleontology

Division Charophyta Migula, 1897
 Class Charophyceae Smith, 1938
 Order Charales Lindley, 1836




Figure 2. Correlation and age of charophyte assemblages. Charophytes and mammalian fossils from the South Gobi area are correlated with the Subeng and Normogen III faunas from the Erlian Basin in Inner Mongolia, China. The age of the South Gobi fossil assemblage is estimated as late Gashatan Asian Land Mammal Age, roughly equivalent to the Clarkforkian North American Land Mammal Age. Modified from Ni et al. (2016).

Family Characeae Agardh, 1824
 Subfamily Charoideae Braun in Migula, 1897
 Genus *Peckichara* Grambast, 1957

Type species.—*Peckichara varians* Grambast, 1957 from the Paleocene to lowermost Eocene (Sparnacian local stage) of the Paris Basin, France.

Peckichara torulosa var. *variens* (Dollfus and Fritel, 1919)
 Sanjuan, Vicente, and Eaton, 2020
 Figure 3.1–3.7

- 1919 *Chara torulosa* Dollfus and Fritel, p. 249, fig. 9.
 1957 *Peckichara torulosa*; Grambast, p. 15, pl. 7, figs. 1–6.
 1957 *Peckichara varians*; Grambast, p. 14, pl. 8, figs. 1–8.
 2020 *Peckichara torulosa* var. *variens*; Sanjuan, Vicente, and Eaton, p. 7, pl. 3, A–F.

Holotype.—Gyrogonites (with collection number not reported) from marls and lignites in Cramant (Marne), Sarron (Oise) according to Dollfus and Fritel (1919, fig. 9).

Occurrence.—*Peckichara torulosa* var. *variens* (originally as *P. varians*) is known from the Paleocene to the early Eocene of Europe and China (Li et al., 2016 and references therein).

Description.—The *P. torulosa* var. *variens* gyrogonites from the South Gobi area are medium to large; 688–770 μm high and 583–708 μm wide. Their isopolarity index ranges between 106 and 118. They are prolate spheroidal to subprolate with seven to eight convolutions in lateral view. The spiral cells are flat to convex, 90–120 μm wide, and ornamented by individualized and prominent tubercles. The tubercles are rounded or oval, usually as wide as the spiral cells, and are spaced regularly (Fig. 3.3, 3.5) or irregularly (Fig. 3.4). The surface ornamentation usually disappears near the apex. The apex is rounded to flat, with a slight to moderate thinning and narrowing of the periapical spiral cells. Sometimes well-developed apical nodules are present, and they can vary in size and shape. Some specimens display small, individualized nodules, and others exhibit thick and large nodules, forming an apical rosette. The base is rounded, with a pentagonal funnel and a pentagonal pore (50–60 μm).




Figure 3. Scanning electron microscope images of fossil charophyte gyrogonites from the latest Paleocene South Gobi area in the Junggar Basin. (1–7) *P. torulosa* var. *variens*: (1, 2) apical view; (3–5) lateral view; (6, 7) basal view. (8–28) *L. vectensis*: (8–14) apical view; (15–21) lateral view; (22–28) basal view. (29–46) *G. lemani capitata*: (29–34) apical view; (35–40) lateral view; (41–46) basal view.

Materials.—Approximately 10 gyrogonites have been recovered from the South Gobi area.

Remarks.—Two varieties of *P. torulosa* have been proposed by Sanjuan et al. (2020). The gyrogonites that bear a mid-cellular crest and massive apical nodules have been termed *P. torulosa* var. *torulosa*, and those that exhibit isolated or irregularly connected tubercles along the spiral

cells and isolated small apical nodules are called *P. torulosa* var. *variens*. The fossil population in this study is referred to *P. torulosa* var. *variens* due to the presence of individualized, prominent, irregularly rounded or elongated tubercles on the spiral cells.

Genus *Lychnothamnus* (Ruprecht, 1845) von Leonhardi, 1863 emend. Braun in Braun and Nordstedt, 1882




Figure 4. Morphometric data of *Lychnothamnus vectensis* (30 gyrogonites) in the South Gobi area, Junggar Basin. (1–4) Frequency distributions: (1) gyrogonite height; (2) gyrogonite width; (3) isopolarity index; (4) number of convolutions. (5) Dispersion graph of gyrogonite height/gyrogonite width.

Type species.—*Lychnothamnus barbatus* (originally as *Chara barbata*) Meyen, 1827 from the Cenozoic of “Plötzensee bei Berlin” (Germany).

Lychnothamnus vectensis (Groves, 1926) Soulié-Märsche, 1989
Figures 3.8–3.28, 4

1926 *Chara vectensis* Groves, p. 172, pl. 2, figs. 2–8.

1958 *Stephanochara vectensis*; Grambast, p. 158.

1989 *Lychnothamnus vectensis*; Soulié-Märsche, p. 160.

Holotype.—Gyrogonites (V. 18321 and V. 18355) from the Bembridge beds in Gurnard Bay on the Isle of Wight, UK (Groves, 1926, pl. 2, figs. 2–8).

Occurrence.—*Lychnothamnus vectensis* was reported first in the late Eocene Bembridge beds in Gurnard Bay on the Isle of Wight, UK (Groves, 1926). It was reported later from the late Eocene of the Paris Basin (Riveline, 1986) and the latest Priabonian to earliest Rupelian of the Ebro Basin (Sanjuan and Martín-Closas, 2014). In the United States, Vicente et al. (2020) and Sanjuan et al. (2020) reported *Lychnothamnus* sp. (probably a form with an affinity to *L. vectensis*) in the early Eocene of Utah. In China, *L. vectensis* is known from the Eocene of the Qaidam Basin on the Tibetan Plateau (Li et al., 2020), and *Lychnothamnus* aff. *L. vectensis* was reported from the Songliao Basin of China by Li et al. (2019).

Description.—The *Lychnothamnus vectensis* gyrogonites from the latest Paleocene South Gobi area are medium to large; 638–845 µm high (mean: 737 µm; standard deviation: 50) and 531–660 µm wide (mean: 596 µm; standard deviation: 32). The isopolarity index ranges from 114 to 149 (mean: 124; standard deviation: 8). They are prolate spheroidal to subprolate with 8–10 convolutions visible in lateral view. The spiral cells are generally convex, but also sometimes flat, and their width ranges between 75 and 100 µm. The apex of the periapical spiral cell is rounded and displays marked thinning. Apical nodules usually are present, and sometimes the nodules are comma or crescent shaped. The apical rosettes are usually well developed, with the apical nodules joined completely at the apical center to form a highly protruding cap. The base is rounded, with a funnel and a pentagonal basal pore ~30–40 µm.

Materials.—Thirty gyrogonites have been recovered from the South Gobi area.

Remarks.—The size of the gyrogonites of *L. vectensis* is normal for each population measured separately (Sanjuan and Martín-Closas, 2015b). However, the gyrogonite’s height range becomes extremely large if all polymorphs from different localities and different latitudes are considered together. Sanjuan and Martín-Closas (2015b) showed that the variation in size of the gyrogonites exhibits a latitudinal polarity according to the morphometric analysis carried out in three populations of *L. vectensis* from the Hampshire, Paris, and Ebro Basins in Europe. The size of the South Gobi population (current latitude: 46°5′19″N) is similar to that of the Paris Basin (current latitude: ~48°N).

In contrast to the type *L. vectensis* population from the Isle of Wight (Groves, 1926; Sanjuan and Martín-Closas, 2015b), with more prominent apical rosette and apical nodules being separated, the apical nodules of *Lychnothamnus* aff. *L. vectensis* in the Songliao Basin population are fused. However, the ornamentation of gyrogonites may change over time, considering the evolutionary trends of other well-supported charophyte anagenetic lineages (e.g., in lineage *Peckichara pectinata* Grambast, 1971; Vicente and Martín-Closas, 2018). *Lychnothamnus* aff. *L. vectensis* probably represents an ancestor of *L. vectensis* s.s.

Genus *Gyrogona* Lamarck, 1804 ex Lamarck, 1822 emend.
Grambast, 1956

Type species.—*Gyrogona medicaginula* Larmark, 1804 from the Oligocene cherts of the Paris Basin, France.

Gyrogona lemani capitata Grambast and
Grambast-Fessard, 1981
Figures 3.29–3.46, 5

1822 *Chara lemani* Brongniart, p. 322, pl. 17, fig. 4.

1927 *Gyrogonites lemani*; Pia, p. 90.

1981 *Gyrogona lemani capitata*; Grambast and Grambast-Fessard, p. 12, pl. 2, figs. 8–12.

Holotype.—Gyrogonites (C121-7) from the Pont-Bernard, commune de Brény (Aisne), France (Grambast and Grambast-Fessard, 1981, p. 12, pl. 2, figs. 8–12).




Figure 5. Morphometric data of *Gyrogonia lemani capitata* (99 gyrogonites) in the South Gobi area, Junggar Basin. (1–4) Frequency distributions: (1) gyrogonite height; (2) gyrogonite width; (3) isopolarity index; (4) number of convolutions. (5) Dispersion graph of gyrogonite height/gyrogonite width.

Occurrence.—*Gyrogonia lemani capitata* was reported in the Lutetian of France, including Pont-Bernard, commune de Brény (Aisne), Cuis (Marne), Pargnan, and Neuilly-Saint-Front (Grambast and Grambast-Fessard, 1981).

Description.—The *G. lemani capitata* gyrogonites are large, with a height of 701–892 µm (mean: 784 µm; standard deviation: 38) and a width of 634–838 µm (mean: 739 µm; standard deviation: 37). Their isopolarity index ranges between 96 and 119 (mean: 106; standard deviation: 5). The shape is oblate spheroidal to prolate spheroidal, with seven to eight convolutions visible in lateral view. The spiral cells are flat to convex, with a width of 115–130 µm. The apex of the gyrogonite is rounded, with moderate periapical thinning of the spiral cells, and ornamented with apical nodules on convex specimens. The base is rounded, with a pentagonal basal pore ~30–40 µm in diameter.

Materials.—More than 100 gyrogonites have been recovered from the South Gobi area.

Remarks.—The *G. lemani capitata* population in this study is smaller in size than that from Pont-Bernard, commune de Brény (Aisne) in Grambast and Grambast-Fessard (1981), which is 950–1,250 µm high and 925–1,200 µm wide. However, the isopolarity index and number of convolutions in

the two populations are similar, with an isopolarity index of 96–116 and six to eight convolutions for the French population.

Discussion

The late Paleocene fossil charophytes recorded in the South Gobi area include the oldest global record of *Gyrogonia lemani capitata* and the second-oldest record for *Lychnothamnus vectensis*. The latest Paleocene *G. lemani capitata* from the South Gobi area is older than the middle Eocene record from France (Fig. 6). The oldest record of *L. vectensis* (as *Lychnothamnus* aff. *L. vectensis*) is from the Cretaceous–Paleocene transition deposits of the Songliao Basin in northeastern China (Li et al., 2019), and the oldest European record is from the late Eocene–early Oligocene transition (Fig. 6). In the United States, its oldest record is probably early Eocene (Vicente et al., 2020). These species clearly were present in Asia for some time before their dispersal to North America and Europe. The older age of the American *Lychnothamnus* aff. *L. vectensis* relative to those in Europe could suggest a pattern of dispersal from Asia to North America first. The dispersal of *L. vectensis* to Europe in the late Eocene could have occurred alongside known tetrapod dispersal events between Europe and North America or through a route (according to known mammalian dispersal patterns) from Asia to Europe (Stidham and Ni, 2014; Stidham et al., 2020). The occurrence of these widespread charophyte taxa in Asia (and North America) before their first records in Europe parallels that of a widely distributed bird group known in the middle Eocene of North America (Utah) and Central Asia (Uzbekistan) before their oldest record in the late Eocene of Western Europe (Stidham et al., 2020). For the Asia-to-Europe potential dispersal route, given physical distances and known geographic barriers (i.e., the Turgai Strait and Tethys Seaway), the charophyte lineages likely would have dispersed from Central Asia to Eastern Europe across the straits in the middle to late Eocene.

Biotic factors could have led to the intercontinental dispersal of charophytes. Birds are known to carry charophyte oospores in their stomachs and externally in mud adhering to their feet and feathers (Proctor, 1962, 1980; Proctor et al., 1967; Trabelsi et al., 2015). Through the alimentary tract of common migratory water birds, many freshwater algae probably could be carried readily between lakes 100–150 miles apart, and through the caecum, they may be carried several times this distance (Proctor, 1959). For example, oospores were found in the digestive tracts of 24% of the birds shot in the Camargue, France (including the red-crested pochard, *Netta rufina* Pallas, 1773), with charophytes, including oospores and vegetative parts, representing 21% of the stomach contents of the ducks (Allouche et al., 1988). Eurasian teal (*Anas crecca* Linnaeus, 1758) also ingested propagules of charophyte macrophytes, with 23% of the total weight ingested consisting of charophyte oospores (Tamisier, 1971). Furthermore, Proctor (1962) found that 34% of the oospores taken from mallards (*Anas platyrhynchos* Linnaeus, 1758) were able to germinate.

Our results show the direction and timing of charophyte dispersal is penecontemporaneous with the intercontinental dispersals of some extinct birds. Those bird records include a group of small-sized pangalliforms in the middle Eocene of Central Asia, Europe, and Africa, along with North America (Stidham




Figure 6. Potential biogeographic dispersal routes of *Lychnothmanus vectensis* and *Gyrogona lemani capitata*. *Lychnothmanus vectensis*: its earliest record is ~300 kyr before the Cretaceous–Paleogene boundary transition, distributed in Xinjiang in the latest Paleocene, in the early Eocene of the United States, and in Spain and France in late Eocene–early Oligocene. *Gyrogona lemani capitata*: its earliest record is in the latest Paleocene in Xinjiang, then dispersed to France in the middle Eocene. The different starred localities are numbered chronologically. Paleogeographic map of the Earth at ~60 Ma reproduced with permission from “Global Paleogeography and Tectonics in Deep Time ©2016 Colorado Plateau Geosystems Inc.”

et al., 2020), ameghinornithid-like birds in the middle Eocene of Inner Mongolia, Eocene and Oligocene of Europe, and early Oligocene of Egypt (Stidham and Smith, 2015; Stidham and Wang, 2017 and references therein), and the stem anatids Romainvilliinae from the late Eocene of Xinjiang and the late Eocene and Oligocene of Europe (Stidham and Ni, 2014). The geographic distributions of those birds imply intercontinental dispersals in the Eocene across the Turgai Strait, demonstrating that the dispersal mechanism of charophytes along with birds is plausible. Furthermore, considering the hypothesized dispersals of the lineages of *Lychnothamnus stockmansii*–*L. major* and *Nitellopsis (Tectochara) merianii*–*N. obtusa* from Europe to Asia in the late Eocene (Sanjuan and Martín-Closas, 2015b), the dispersal routes of charophytes were likely bidirectional in the Eocene between Europe and Asia.

We explored the sedimentary facies of Asia, North America, and Europe to interpret the paleoenvironments of charophytes when they arrived. Proctor (1959) mentioned suitable media or environmental conditions are important for *Chara* zygotes to germinate and colonize. *Gyrogona lemani capitata* is found in sandstones and sandy mudstones attributed to deposition in intermittent riverine environments in the uppermost Paleocene of the northeastern Junggar Basin, Xinjiang (Ni et al., 2016). It was later found in middle Eocene beds of France. It occurs in marls from Pont-Bernard, commune de Brény (Aisne), Cuis (Marne), Pargnan, and Neuilly-Saint-Front of France, corresponding to sedimentation in shallow lacustrine facies (Grambast and Grambast-Fessard, 1981). Therefore, it appears that either *G. lemani capitata* shifted or varied its habitat preferences during its biogeographic history, from fluvial environments in the South Gobi area to shallow lacustrine belts in the middle Eocene of Europe, or it occupied a wider array of habitats

than previously recognized. *Lychnothmanus* aff. *L. vectensis* was reported from shallow lacustrine facies of the Songliao Basin (Li et al., 2019) and in lake delta front facies of the Jiaolai Basin, eastern China. To the west, in the northeastern Junggar Basin, Xinjiang, *Lychnothmanus* aff. *L. vectensis* is present in fluvial sandstones and sandy mudstones. *Lychnothmanus* aff. *L. vectensis* recovered from Utah (United States) came from beds of siltstones containing mainly thin thalli, ostracods, gastropods, and trace fossils that were interpreted as representing floodplain environments. Assuming all these findings represent autochthonous populations, these data suggest the species occupied a wide range of habitats in China. When it dispersed to Europe around the Eocene–Oligocene boundary transition, *L. vectensis* appears to have become restricted in its habitats to shallow lacustrine facies. For example, *L. vectensis* is in the shallow lacustrine deposits of the Artés Formation, Ebro Basin, Catalonia (Sanjuan and Martín-Closas, 2015b) and the lacustrine bluish marls of the Argenteuil Formation in Loiret (Centre-Val de Loire) in the Paris Basin, France (Riveline, 1986). In addition, *L. vectensis* was found in the lacustrine deposits from the Bembridge Marls Member (Bouldnor Formation, Solent Group) in the Hampshire Basin on the Isle of Wight, United Kingdom (Sanjuan and Martín-Closas, 2015b and references therein).

Conclusions

We studied the taxonomy, paleoecology, and paleobiogeography of a latest Paleocene charophyte flora in the South Gobi area of Xinjiang in China, which includes the taxa *Peckichara torulosa* var. *varians*, *Lychnothmanus vectensis*, and *Gyrogona lemani capitata*. The dispersal of this flora from Asia to Europe in the middle to late Eocene probably occurred before the retreat

of the Turgai Strait in both the Tarim area and the Siberian Basin by the end of the late Eocene and before the “Grande Coupure” in Europe and the “Mongolian Remodelling” event in Asia during the Eocene–Oligocene transition. Waterbirds may have played an important role in facilitating this intercontinental dispersal. In comparison with European Eocene records, it appears that *G. lemani capitata* and *Lychnothamnus* aff. *L. vectensis* had wider paleoecological tolerances in the Paleocene of China, before their migration westward.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (grant numbers 41625005, 41972018, 41730317, 41888101, 41988101, 41902020), the Strategic Priority Research Program of Chinese Academy of Sciences (grant numbers CAS XDB26030400, XDB26030300, XDB26010304, XDA20070203, XDA19050100), and the open project of State Key Laboratory of Paleobiology and Stratigraphy (grant number 193103). We thank G. Wang, R. Li, and Y. Wu for sorting the specimens from the screen-washing concentrates. Thanks to X. Jin for help with SEM imaging. We gratefully acknowledge C. Martín-Closas (University of Barcelona) and an anonymous reviewer for their careful work and thoughtful suggestions that have helped improve this paper substantially.

References

- Agardh, C.A., 1824, *Systema Algarum*: Lundae, Literis Berlingianis, 312 p.
- Allouche, L., Roux, P., and Tamisier, A., 1988, Position trophique des nettes rousses (*Netta rufina*, Pallas) hivernant en Camargue: *Revue d'écologie-la terre et la vie*, v. 43, p. 167–175.
- Bhatia, S.B., 2006, Ecological parameters and dispersal routes of *Lychnothamnus barbatus* (Characeae) in the early–middle Holocene from the Ganga plain, India, *Cryptogamie: Algologie*, v. 27, p. 341–356.
- Bosboom, R., Mandic, O., Dupont-Nivet, G., Proust, J.-N., Ormukov, C., and Aminov, J., 2017, Late Eocene palaeogeography of the proto-Paratethys Sea in Central Asia (NW China, southern Kyrgyzstan and SW Tajikistan): *Geological Society, London, Special Publications*, v. 427, p. 565–588.
- Braun, A., and Nordstedt, C.F.O., 1882, *Fragmente einer Monographie der Characeen*: Berlin, Königlichem Akademie der Wissenschaften, 211 p.
- Brongniart, A., 1822, Sur la classification et la distribution des végétaux fossiles en général, et sur ceux des terrains de sédiment supérieur en particulier: *Mémoires du Muséum d'histoire naturelle Paris*, 91 p.
- Dollfus, G.F., and Fritel, P.H., 1919, Catalogue raisonné des characées fossiles du Bassin de Paris: *Bulletin de la Société Géologique de France*, v. 19, p. 243–261.
- Grambast, L., 1956, Le genre *Gyrogonia* Lmk (Characeae): *Comptes rendus sommaires des Séances de la Société Géologique de France*, v. 14, p. 278–280.
- Grambast, L., 1957, Ornementation de la gyrogonite et systématique chez les Charophytes fossiles: *Revue générale de Botanique*, v. 64, p. 339–362.
- Grambast, L., 1958, Etude sur les Charophytes Tertiaires d'Europe Occidentale et leurs rapports avec les formes actuelles [Ph.D. thesis]: Paris, Paris University, 258 p.
- Grambast, L., 1971, Remarques phylogénétiques et biocronologiques sur les Septorella du Crétacé terminal de Provence et les charophytes associés: *Paléobiologie Cont.*, v. 2, p. 1–38.
- Grambast, L., and Grambast-Fessard, N., 1981, Étude sur les Charophytes tertiaires d'Europe occidentale. III. Le genre *Gyrogonia*: *Paléobiologie Continentale*, v. 12, no. 2, p. 1–35.
- Grambast, L., and Paul, P., 1965, Observations nouvelles sur la flore de charophytes du Stampien du bassin de Paris: *Bulletin de la Société Géologique de France*, v. 7, p. 239–247.
- Grambast, L., and Soulié-Marsche, I., 1972, Sur l'ancienneté et la diversification des *Nitellopsis* (Charophytes): *Paléobiologie Continentale*, v. 3, p. 1–14.
- Groves, J., 1919, Notes on *Lychnothamnus* Braun, *Journal of Botany*, London v. 57, p. 125–129.
- Groves, J., 1926, Charophyta, in Reid, E.M., and Chandler, M.E.J., eds., *Catalogue of Cainozoic plants in the Department of Geology, Volume 1, The Bembridge flora*: London, The British Museum (Natural History), p. 165–173.
- Joyce, W., and Rabi, M., 2015, A revised global biogeography of turtles: *PeerJ Preprints*, <https://doi.org/10.7287/peerj.preprints.853v1>
- Lamarck, J.B., 1804, Suite des mémoires sur les fossiles des environs de Paris: *Annales du Muséum d'histoire naturelle*, v. 5, p. 349–357.
- Lamarck, J.B., 1822, *Histoire Naturelle des Animaux sans Vertèbres*: Paris, Verdère, 711 p.
- Li, S., Wang, Q.F., Zhang, H.C., Lu, H.N., and Martín-Closas, C., 2016, Charophytes from the Cretaceous–Paleogene transition in the Pingyi Basin (eastern China) and their Eurasian correlation: *Cretaceous Research*, v. 59, p. 179–200.
- Li, S., Wang, Q.F., Zhang, H.C., Wan, X.Q., and Martín-Closas, C., 2019, Charophytes from the Cretaceous–Paleocene boundary in the Songliao Basin (north-eastern China): a Chinese biozonation and its calibration to the Geomagnetic Polarity Time Scale: *Papers in Palaeontology*, v. 5, p. 47–81.
- Li, S., Wang, Q., Cui, Q., Zhang, S., Wang, H., and Zhang, H., 2020, Eocene charophyte flora from the Xia Ganchaigou Formation in Zongmahai Lake area, Qaidam Basin, northwest China: *Acta Palaeontologica Sinica*, v. 59, p. 327–337. [in Chinese with English abstract]
- Lindley, J., 1836, *A Natural System of Botany, or, a systematic view of the organization, natural affinities, and geographical distribution, of the whole vegetable kingdom: together with the uses of the most important species in medicine, the arts, and rural or domestic economy*: London, Longman, Rees, Orme, Brown, Green, and Longman, 558 p.
- Linnaeus, C., 1758, *Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis* (tenth edition), v. 1: Stockholm, L. Salvii, 823 p.
- Loiseleur Deslongschamps, J.L.A., 1810, Notice sur les plantes à ajouter à la flore de France (Flora Gallica) avec quelques corrections et observations: Paris, J.B. Sajou, 172 p.
- Martin, T., and Averianov, A.O., 2004, A new docodont (Mammalia) from the Middle Jurassic of Kyrgyzstan, Central Asia: *Journal of Vertebrate Paleontology*, v. 24, p. 195–201.
- Martín-Closas, C., and Wang, Q.F., 2008, Historical biogeography of the lineage *Atopochara trivolvris* Peck 1941 (Cretaceous Charophyta): *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 260, p. 435–451.
- Matthew, W.D., and Granger, W., 1925, Fauna and correlation of the Gashato Formation of Mongolia: *American Museum Novitates*, v. 189, p. 1–12.
- McKenna, M.C., 1975, Fossil mammals and early Eocene North Atlantic Land Continuity: *Annals of the Missouri Botanical Garden*, v. 62, p. 335–353.
- Meng, J., and Wyss, A.R., 2001, The morphology of *Tribosphenomys* (Rodentiaformes, Mammalia): phylogenetic implications for basal Glires: *Journal of Mammalian Evolution*, v. 8, p. 1–71.
- Meng, J., Ni, X., Li, C., Beard, K.C., Gebro, D.L., Wang, Y., and Wang, H., 2007, New material of Alagomyidae (Mammalia, Glires) from the late Paleocene Subeng locality, inner Mongolia: *American Museum Novitates*, v. 3597, p. 1–29.
- Meyen, F.J.F., 1827, Beobachtungen und Bemerkungen über die Gattung *Chara*: *Linnaea*, v. 2, p. 55–81.
- Migula, W., 1897, Die Characeen, in Rabenhorst, G.L., ed., *Kryptogamen-flora von Deutschland, Österreichs und der Schweiz*: Leipzig, Edword Kummer, p. 1890–1897.
- Missiaen, P., and Smith, T., 2005, A new nyctitheriid insectivore from Inner Mongolia (China) and its implications for the origin of the Asian nyctitheriids: *Acta Palaeontologica Polonica*, v. 50, p. 513–522.
- Missiaen, P., and Smith, T., 2008, The Gashatan (late Paleocene) mammal fauna from Subeng, Inner Mongolia, China: *Acta Palaeontologica Polonica*, v. 53, p. 357–378.
- Ni, X., Li, Q., Stidham, T.A., Li, L., Lu, X., and Meng, J., 2016, A late Paleocene probable metatherian (?deltatheroidan) survivor of the Cretaceous mass extinction: *Scientific Reports*, v. 6, art. 38547.
- Pallas, P.S., 1773, *Reise durch verschiedene Provinzen des russischen Reichs*: St. Petersburg, Gedruckt bey Kayserlichen Academie der Wissenschaften.
- Peck, R.E., 1941, Lower Cretaceous Rocky Mountain non marine microfossils: *Journal of Paleontology*, v. 15, p. 285–304.
- Pia, J., 1927, *Thallophyta*, in Hirmer, M., ed., *Handbuch der Paläobotanik*, v. 1: München, R. Oldenbourg Druck und Verlag, p. 31–136.
- Proctor, V.W., 1959, Dispersal of fresh-water algae by migratory water birds: *Science*, v. 130, p. 623–624.
- Proctor, V.W., 1962, Viability of *Chara* oospores taken from migratory water birds: *Ecology*, v. 43, p. 528–529.
- Proctor, V.W., 1980, Historical biogeography of *Chara* (Charophyta): an appraisal of the braun-wood classification plus a falsifiable alternative for future consideration: *Journal of Phycology*, v. 16, p. 218–233.

- Proctor, V.W., Malone, C.R., and DeVlaming, V.L., 1967, Dispersal of aquatic organisms: viability of disseminules recovered from the intestinal tract of captive killdeer: *Ecology*, v. 48, p. 672–676.
- Riveline, J., 1986, Les Charophytes du Paléogène et du Miocène Inférieur d'Europe Occidentale: Paris, Centre National de la Recherche Scientifique, 304 p.
- Ruprecht, F.J., 1845, *Distributio Cryptogarum Vascularum in Imperio Rossico*, Beiträge zur Pflanzenkunde des Russischen Reiches, fasc. 3: St. Petersburg, Kaiserlichen Academie der Wissenschaften, 56 p.
- Sanjuan, J., and Martín-Closas, C., 2014, Taxonomy and palaeobiogeography of charophytes from the upper Eocene–lower Oligocene of the Eastern Ebro Basin (Catalonia, NE Spain): *Geodiversitas*, v. 36, p. 385–420.
- Sanjuan, J., and Martín-Closas, C., 2015a, Biogeographic history of two Eurasian Cenozoic charophyte lineages: *Aquatic Botany*, v. 120, p. 18–30.
- Sanjuan, J., and Martín-Closas, C., 2015b, Gyrogonite polymorphism in two European charophyte biozone index species: *Papers in Palaeontology*, v. 1, p. 1–14.
- Sanjuan, J., Vicente, A., and Eaton, J.G., 2020, New charophyte flora from the Pine Hollow and Claron formations (southwestern Utah). Taxonomic, biostratigraphic, and paleobiogeographic implications: *Review of Palaeobotany and Palynology*, v. 282, art. 104289.
- Smith, G.M., 1938, *Algae and Fungi: Charophyceae*. *Cryptogamic Botany*, v. 1: New York, McGraw Hill, 545 p.
- Smith, T., Van Itterbeeck, J., and Missiaen, P., 2004, Oldest plesiadapiform (Mammalia, Proprimates) of Asia and its paleobiogeographical implications for faunal interchange with North America: *Comptes Rendus Palevol*, v. 3, p. 43–52.
- Soulié-Marsche, I., 1989, Etude comparée de gyrogonites de charophytes actuelles et fossiles et phylogénie des genres actuels: Millau, France, Imprimerie des Tilleuls, 237 p.
- Stidham, T.A., and Ni, X., 2014, Large anseriform (Aves: Anseriformes: Romainvilliinae?) fossils from the late Eocene of Xinjiang, China: *Vertebrata Palasiatica*, v. 52, p. 98–111.
- Stidham, T.A., and Smith, N.A., 2015, An ameghinornithid-like bird (Aves, Cariamae, ?Ameghinornithidae) from the early Oligocene of Egypt: *Palaeontologia Electronica*, v. 18.1.5A, 8 p., <https://palaeo-electronica.org/content/pdfs/470.pdf>
- Stidham, T.A., and Wang, Y., 2017, An ameghinornithid-like bird (Aves: Cariamae: Ameghinornithidae?) from the middle Eocene of Nei Mongol, China: *Vertebrata Palasiatica*, v. 55, p. 218–226.
- Stidham, T.A., Townsend, K.E., and Holroyd, P.A., 2020, Evidence for wide dispersal in a stem galliform clade from a new small-sized middle Eocene pangalliform (Aves: Paraortygidae) from the Uinta Basin of Utah (USA): *Diversity*, v. 12, article 90.
- Tamisier, A., 1971, Régime alimentaire des sarcelles d'hiver *Anas crecca* en Camargue: *Alauda*, v. 39, p. 261–311.
- Tang, W., Zhang, Y., Pe-Piper, G., Piper, D.J.W., Guo, Z., and Li, W., 2020, Soft-sediment deformation structures in alkaline lake deposits of lower Permian Fengcheng Formation, Junggar Basin, NW China: implications for syn-sedimentary tectonic activity: *Sedimentary Geology*, v. 406, art. 105719.
- Trabelsi, K., Sames, B., Salmouna, A., Piovesan, E. K., Rouina, S.B., Houla, Y., Touis, J., and Soussi, M., 2015, Ostracods from the marginal coastal Lower Cretaceous (Aptian) of the Central Tunisian Atlas (North Africa): paleo-environment, biostratigraphy and paleobiogeography: *Revue de Micropaléontologie*, v. 58, p. 309–331.
- Unger, F., 1852, *Iconographia plantarum fossilium: Denkschriften Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse*, Wien, v. 4, p. 73–118.
- Vicente, A., and Martín-Closas, C., 2018, Gradualistic characean lineages in the Upper Cretaceous–Palaeocene of Southern Europe: *Historical Biology*, v. 30, p. 593–607.
- Vicente, A., Sanjuan, J., Eaton, J.G., and Villanueva-Amadoz, U., 2020, The oldest record of North American *Lychnothamnus* (northeastern Sonora, Mexico): implications for the evolution, ecology, and paleogeographic distribution of the genus: *Aquatic Botany*, v. 167, art. 103271.
- von Leonhardi, H.F., 1863, Über die böhmischen Characeen: *Lotos*, v. 13, p. 55–62.
- Wang, Y.Q., Meng, J., Ni, X.J., and Beard, K.C., 2008, A new early Eocene arctostylopoid (Arctostylopida, Mammalia) from the Erlan Basin, Nei Mongol (Inner Mongolia), China: *Journal of Vertebrate Paleontology*, v. 28, p. 553–558.

Accepted: 25 November 2021